POSSIBLE OPIOID OVERDOSE: WHAT TO DO

The person might have used medications or illicit drugs and shows the following signs:



No response to sound or pain



Laboured or snore-like breathing, or absence of breathing

The person seems to be unconscious

A POSSIBLE OVERDOSE

TRY MAKING THEM
RESPOND TO SOUND
OR PAIN



YELL their name
TALK TO THEM loudly

RUB the centre of their chest (sternum) hard


They are unresponsive

RECOGNIZING

CALL OR HAVE
SOMEONE
CALL
911

IF YOU ARE ON YOUR OWN WITHOUT A PHONE:

- Inject a dose of naloxone
- Perform chest compressions for 2 min
- Lay the person on their side
- Find a way to call 911 and follow instructions



They are unresponsive

INJECT
A DOSE
OF NALOXONE











Inject naloxone into thigh or shoulder muscle (at a 90° angle)



They are unresponsive

PERFORM
CHEST COMPRESSIONS
IMMEDIATELY



GIVE 2 compressions (5 cm deep) per second

OR

START giving CPR if you have been trained, using the barrier mask

They are unresponsive **3 min** after injection

INJECT
ANOTHER DOSE
OF NALOXONE



REPEAT STEPS 4 AND 5

as long as the person is unresponsive



Out of naloxone?

Continue chest compressions or CPR until help arrives

IF THE PERSON WAKES UP...

Lay them on their side. Explain what just happened and how it's important that they be seen by a health professional. The risk of another overdose after the first one is high because naloxone has a short duration of action.

Tell the person that they should not use opioids in the next few hours to avoid another overdose.

Stay with them while waiting for help. The Good Samaritan Law provides protection for overdose witnesses and victims.



- · Store naloxone in a cool, dark place
- Check the expiry date
- Do not fill the syringes in advance
- Naloxone can be injected through light clothing
- Dispose of used syringes and needles safely

For more information, visit Québec.ca/opioides

Produced by the Institut national d'excellence en santé et en services sociaux





233-01A © Gouvernement du Québec, 2018