

၎င်းကဏ္ဍအတွက် အောက်ဖော်ပြပါအတိုင်း ဖော်ပြပါမူဝါဒများကို အသုံးပြုနိုင်ပါသည်။

[illegible][illegible]

- Babies born during the RSV season.
- Babies under 8 months of age at the start of
- RSV season who have not yet received treatment.
- Babies under 19 months who have a condition that increases their risk of severe RSV infection.

- [illegible]

The preventive treatment is safe. In most cases, it will not cause a reaction. Check with your CLSC, a health professional or your midwife at the time of your next visit.

ԿՆՉԴԼԾՍՈՒ ՎՇՈՔՈՂԻՄ. ՎԴՐՏԻՏԻՄ,
ԴԿՆԻՍՈՒԴՆԻՄՈՂԻՄ:

[illegible]

The RSV preventive treatment is offered during the RSV season which is usually in the fall and winter. It is offered to all newborns before they leave the hospital or birth centre. If your baby did not get it at birth, they may get it at a local point of service or a local community services center (CLSC).

[illegible]

- Wash your hands often.
- Cough or sneeze into your elbow.
- Stay home if you have symptoms.

- ልዩጋዬዎዊርቲኬርጋር ልህሰር.
- ነፃጋርጋር ርዕድጋርጋር ልዩጋዬዎዊርቲኬር ልህሰር.
- ልዩጋዬዎዊርቲኬር ልህሰርጋርጋር